

苏丹 IV 酒精饱和溶液

货号: G1521

规格: 100mL

保存: 室温, 避光保存, 有效期 1 年。

产品介绍:

脂质(Lipid)是中性脂肪、类脂及其衍生物的总称,其共同的物理特性是不溶于水,易溶于有机溶剂(如乙醇、乙醚等)。中性脂肪(Neutral fat)是由三分子脂肪酸和一分子甘油组成的脂类,呈中性。中性脂肪是储存能量的方式之一,在氧化时释放出能量。在正常情况下,除脂肪细胞外,其他细胞在光学显微镜下几乎看不到脂滴,如果细胞质内出现大量脂滴即为脂肪变性,常见于肝细胞、心肌细胞、肾曲管上皮细胞等。中性脂肪染色经常采用苏丹 II、苏丹 III、苏丹 IV、苏丹黑 B、油红 O 法等。苏丹染料脂质染色的机理一般认为纯属物理学的脂溶作用和吸附作用。苏丹类染料由于在脂质中的溶解度大于在有机溶剂的溶解度,所以染色时染料便从染液中转移到被染的脂质中去,使脂质呈现出染液的颜色。

苏丹 IV 酒精饱和溶液主要用于显示组织器官的脂肪变性和类脂质的异常沉着,常发生于肝、肾、心等实质脏器的脂肪变性,细胞内出现多数中性脂肪滴;鉴别和诊断脂肪组织中所发生的肿瘤及其性质。标本不采用含有乙醇的固定液(如需要固定可采用 10%福尔马林)、也不采用石蜡切片,需用冰冻切片或碳蜡切片。

自备材料:

载玻片、70%乙醇、蒸馏水

操作步骤: (仅供参考)

1. 新鲜组织低温切片,一般-20℃到-25℃。如样本为脂肪瘤,应调节至-30℃。
2. 冰冻切片 5-10 μ m(6-8 μ m 为佳),贴于载玻片上。70%乙醇稍微浸洗一下。
3. 入苏丹 IV 酒精饱和溶液,浸染 1-2min。
4. 70%乙醇洗去多余染液。蒸馏水浸洗 1min。
5. 按实验具体要求进行下游实验。

染色结果:

中性脂肪	猩红色
------	-----

注意事项:

1. 脂肪染色时标本不宜采用含有乙醇的固定液、也不宜用石蜡切片,需用冰冻切片。
2. 在染色过程中必须防止染料发生沉淀。故切片入染液时应密封,勿与流动空气相接触,避免溶液挥发时发生沉淀。
3. 冰冻切片较易着色,复染时应避免过染。
4. 苏丹染料容易褪色,应密闭保存。
5. 为了您的安全和健康,请穿实验服并戴一次性手套操作。

Sudan IV Saturated Solution, Ethanol Solvent

Cat: G1521

Size: 100mL

Storage: RT, avoid light, valid for 1 year.

Introduction

Neutral fat stains often use Sudan II, Sudan III, Sudan IV, Sudan black B, oil red O and so on. The dyeing principle is generally regarded as a physical miscibility or adsorption, and the fat is dyed by miscibility. The solubility of dyes in frozen sections is higher than that in the original solvent, so when dyeing, the dyes are transferred from organic solvents to lipids and the fats are dyed.

Sudan IV Saturated Solution, Ethanol Solvent is mainly used to show fatty degeneration of tissues and organs and abnormal lipid-like sedation. It often occurs in fatty degeneration of liver, kidney, heart and other parenchymal organs that most of the neutral fat droplets appear in cells. It can identify and diagnose tumors and their properties in adipose tissue. The sample should not fixed with ethanol (10% formalin if required) or paraffin. Frozen sections or carbon wax sections were needed.

Self Provided Materials

Slide, 70% ethanol, Distilled Water

Protocol (for reference only)

1. Prepare low-temperature sections of fresh tissue, generally at -20°C to -25°C . If the sample is lipoma, it should be adjusted to -30°C .
2. Slice the frozen section at 5-10 micron (6-8 micron is preferable) and attach to the slide.
3. Wash slightly with 70% ethanol.
4. Stain with Sudan IV Saturated Solution, Ethanol Solvent for 1-2min.
5. Wash with 70% alcohol to remove excess solution. Rinse with distilled water for 1 min.
6. Follow the next steps according to the specific requirements of the experiment.

Result

Neutral fat	Scarlet
-------------	---------

Note

1. Samples should not be fixed with ethanol or embedded with paraffin, and frozen sections should be used. It is recommended to use 10% neutral formalin or 10% formaldehyde-calcium solution to fix.
2. Precipitation of dyes must be prevented during dyeing. Therefore, slices should be covered when they are put into dye solution, so as not to contact with flowing air to avoid precipitation when the solution volatilizes.
3. Frozen sections are easier to be colored, and over-dyeing should be avoided when re-dyeing with Mayer hematoxylin.
4. Sudanese dyes fade easily and should be kept in airtight condition.
5. For your safety and health, please wear lab clothes and disposable gloves.