

Soil FDA Hydrolase Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment: Spectrophotometer

Cat No: BC0480

Size:50T/24S

Components:

Reagent I: 30 mL×1, storage at 4°C.

Reagent II: Powder×1, storage at -20°C and protect from light. Add 3 mL of acetone to dissolve before use.

Standard: Powder×1, 10 mg of fluorescein, storage at -20°C and protect from light. Before use, add 3.03 mL of 50% acetone (acetone(V): distilled water(V)=1:1) to prepare 10 μmol/mL fluorescein standard solution, which can be dissolved in a 45°C water bath.

Product Description:

Fluorescein diacetate (FDA) hydrolysis is one of the most important biological indicators in the study of soil quality, which can reflect the activity of soil microbial, the change of soil quality and the transformation of organic matter in ecosystem.

FDA is a colorless compound, which can be hydrolyzed by many soil enzymes in the medium. After dehydration reaction, fluorescein is the final product of enzymatic hydrolysis. The fluorescein is stable and not easy to be decomposed, and has a strong absorption peak at 490 nm. The activity of FDA hydrolase can be calculated by detecting the change of absorption value at 490 nm.

Reagents and Equipment Required but Not Provided:

Balance, low temperature centrifuge, spectrophotometer, 1 mL glass cuvette, constant temperature water bath, acetone, 30 mesh sieve (or smaller).

Procedure:

I. Treatment of soil samples:

Natural air drying of fresh soil sample or air drying in 37°C oven, passing through a 30 mesh sieve.

II. Determination steps

1. Preheat spectrophotometer for 30 minutes, adjust the wavelength to 490 nm, set zero with 50% acetone.
2. Dilute 10 μmol/mL fluorescein standard solution with 50% acetone to 2, 0 μmol/mL standard solution (0 is blank tube). Take 1 mL in the 1 mL glass cuvette to determine the absorbance A_S , A_B at 490 nm separately, calculate $\Delta A_S = A_S - A_B$.
3. Add reagents as the following table.

Reagent name	Control tube (C)	Test tube (T)
Sample (g)	0.1	0.1

Reagent I (μL)	500	500
Acetone (μL)	450	-
Reagent II	50	50
Mix well, shake for 1 hour at 37°C.		
Acetone (μL)	-	450
Centrifuge at 10000 ×g for 5 minutes at 25 °C, take the supernatant, measure the absorbance (A) at 490 nm, recorded as A _T , A _C . ΔA _T = A _T -A _C .		

Note: The blank tube and standard tube only need to be measured once or twice.

III. The calculation formula of FDA hydrolase activity:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the production 1 μmol of fluorescein per day every gram of soil sample.

$$\text{FDA (U/g prot)} = (\Delta A \div \Delta A_S \times C_S) \times V_{RT} \div W \div T = 48 \times (\Delta A \div \Delta A_S) \div W$$

V_{RT}: The total volume of reaction, 1 mL;

C_S: Concentration of standard solution, 2 μmol/mL;

T: Catalytic reaction time, 1 hour = 1/24 day;

W: Weight of air dried sample, g.

Note:

1. Try to use fresh soil samples or samples preserved under short-term low temperature conditions, otherwise it is difficult to accurately reflect the enzyme activity.
2. Carry out the pre-experiment before the determination. If the absorbance value is greater than 1.2, please carry out the determination to reduce the mass of soil sample, and multiply the dilution multiple in the calculation formula. If the absorption value is too small, it can be determined by increasing the mass of soil sample or reaction time.

Experimental example:

1. Take 0.1g of clover soil into 1.5mlep tube, as control tube and measuring tube respectively, operate according to the determination steps, take the supernatant and dilute it 5 times for determination, and calculate $\Delta A = A_T - A_C = 1.148 - 0.066 = 1.082$, $\Delta A_S = A_S - A_B = 0.499 - 0 = 0.499$ according to the soil mass
 FDA activity (U/g soil sample) = $48 \times (\Delta A \div \Delta A_S) \times W \times 5$ (dilution ratio) = $48 \times (1.082 \div 0.499) \times 0.1 \times 5$ (dilution ratio) = 5204.01 U/g soil sample.

2. Take 0.1g of forest soil in 1.5mlep tube, as control tube and determination tube respectively, operate according to the determination steps, take the supernatant diluted 5 times for determination, calculate $\Delta A = A_T - A_C = 0.973 - 0.133 = 0.84$, $\Delta A_S = A_S - A_B = 0.499 - 0 = 0.499$, calculate the enzyme activity according to the soil mass

FDA activity (U/g soil sample) = $48 \times (\Delta A \div \Delta A_S) \times W \times 5$ (dilution ratio) = $48 \times (0.84 \div 0.499) \times 0.1 \times 5$ (dilution ratio) = 4040.08 U/g soil sample.

References:

- [1] Sánchez-Monedero M A, Mondini C, Cayuela M L, et al. Fluorescein diacetate hydrolysis,

respiration and microbial biomass in freshly amended soils[J]. *Biology and Fertility of Soils*, 2008, 44(6): 885-890.

[2] Paudel B R, Udawatta R P, Anderson S H. Agroforestry and grass buffer effects on soil quality parameters for grazed pasture and row-crop systems[J]. *Applied Soil Ecology*, 2011, 48(2): 125-132.

Related Products:

BC0240/BC0245 Soil Saccharase(S-SC) Activity Assay Kit

BC0140/BC0145 Soil Acid Phosphatase(S-ACP) Activity Assay

BC0460/BC0465 Soil Neutral Phosphatase(S-NP) Activity Assay Kit

BC0280/BC0285 Soil Alkaline Phosphatase(S-AKP/ALP) Activity Assay Kit